

Catalogue of American Amphibians and Reptiles.

John, R.R., E.J. Bentz, M.J. Rivera Rodríguez, A.M. Bauer, and R. Powell. 2012. *Bachia heteropa*.

***Bachia heteropa* (Lichtenstein and von Martens)**

Earless Worm Lizard

Chalcides heteropus Lichtenstein and von Martens in Lichtenstein 1856:17. Type-locality, "La Guayra [= La Guaira, Distrito Federal], Venezuela." Lectotype, Zoologisches Museum Berlin (ZMB) 1173 (designated by Bauer and Günther 1994), an adult collected by J. Gollmer, sex and date of collection unknown (examined by AMB). See **Remarks**.

Bachia heteropus: Garman 1892:97.

Bachia heteropa: Beebe 1944:14.

Bachia heretopa: Ávila Pires 2005:32. *Lapsus*.

• **CONTENT.** Five subspecies are currently recognized: *Bachia h. heteropa*, *B. h. alleni*, *B. h. lineata*, *B. h. marcelae*, and *B. h. trinitatis*. Hailey and Caza-bon-Mannette (2011) suggested that the population on Tobago might represent an endemic subspecies. Galis et al. (2010) and Kohlsdorf et al. (2010) both presented phylogenies suggesting that *B. heteropa* as currently defined is polyphyletic. See also **Remarks**.

• **DEFINITION.** *Bachia heteropa* is a small member of the genus *Bachia* (maximum known SVL 64 mm). Body and tail are elongated, limbs are reduced with 4 toes on forelimbs and 2–4 toes on hindlimbs (varying by subspecies), external ears are absent, eye size is reduced, and lower eyelids bear an unsegmented window. Dorsal scales are smooth, hexagonal, and imbricate, in 5–12 longitudinal rows; lateral and ventral scales are smooth, quadrangular, and juxtaposed, in 16–22 longitudinal rows; tail scales are mostly hexagonal (Dixon 1973). Dorsal scales number 37–41 from occiput to vent, scale rows around midbody (SAB) are 25–31, gulars 5–8, ventrals 25–29, supraoculars 2–3, superciliaries 3, supralabials and infralabials usually 6/6 or 6/5, labials and parietals not in contact, interparietal, frontonasal, nasals,



FIGURE 1. *Bachia heteropa alleni* from Friendship Bay, Bequia, St. Vincent and the Grenadines. Photograph by S. Blair Hedges.




FIGURE 2. *Bachia heteropa heteropa* (MHNLS 9207) from El Guacuco, San Juan de Las Galdonas, Península de Paria, Sucre, Venezuela. Drawing by Osvaldo Villareal (courtesy of Gilson Rivas).





FIGURE 3. *Bachia anomala* (= *B. heteropa lineata*) (from Brongersma 1946).

and parietals present, prefrontals absent or present in median contact, second pair of chin shields not usually in medial contact and do not typically reach the oral border, males usually with 2/2 preanal pores (absent in females), and preanal shields usually 4, but 3–5 occurring once in 80 specimens (Dixon 1973; Schwartz and Henderson 1991).

Coloration varies slightly between subspecies, but all have a basic pattern of 3 longitudinal dorsal dark lines alternating with 4 wider, light interspaces extending from the parietals onto the tail. The sides of the body are dark brown to blackish and the venter is light brown to yellowish tan. The upper edge of the lateral dark stripe tends to be darker than the stripe. In all subspecies, giving an illusion of 5 dark stripes on the body (Dixon 1973).

• **DIAGNOSIS.** *Bachia heteropa* can be distinguished from sympatric South American congeners by scale pattern. Dorsal scales in the *B. heteropa* group (*B. heteropa* and *B. pallidiceps*) are hexagonal, smooth, and imbricate, and lateral and ventral scales



MAP. Distribution of *Bachia heteropa*; circles mark type-localities and dots indicate other known locality records. Note the type-locality of *Chalcides heteropus* (La Guayra [= La Guaira, Distrito Federal], Venezuela), marked with a circle and an "x," falls within the range of *B. heteropa lineata* according to the ranges shown by Dixon (1973). However, some authors (e.g., Rivas Funemayor et al. 2005) called into question the validity of the various subspecies, citing clinal variation across the range of the species. Gollmer's collections were probably derived from a wider area within Venezuela than just La Guaira, and that locality may merely indicate the point from which the specimens were shipped. Lichtenstein was notorious for taking collectors' detailed data and replacing them with broad geographic reference points. Uncertain records from Colombia are not shown (see **Remarks** and **Comment**).




FIGURE 4. Progressive reduction in size and loss of prefrontal scales and hindlimb toes in *Bachia heteropa* from more mesic habitats in the eastern portion of the species' distribution to increasingly xeric habitats in the west: (A) *B. h. alleni*, (B) *B. h. trinitatis* from Trinidad, (C) *B. h. trinitatis* from the mainland, (D) *B. h. heteropa*, and (E) *B. h. lineata* (from Dixon 1973).

are quadrangular, smooth and juxtaposed; species in the *B. bresslaui* group have keeled hexagonal scales dorsally as well as laterally; species in the *B. dobignyi* group also have hexagonal scales dorsally and laterally, but these are smooth; species in the *B. flav-*

escens group have quadrangular scales all over the body (no hexagonal scales). *Bachia heteropa* can be distinguished from *B. pallidiceps* in having 3 superciliaries and 24+ (mean 28.4) midbody scale rows, whereas *B. pallidiceps* has 2 superciliaries and ≤ 24

(mean 22.9) midbody scale rows (Dixon 1973). *Bachia heteropa* can be distinguished from all sympatric Lesser Antillean lizards by the following combination of characters (Schwartz and Henderson 1985): large, square, ventral plates, fewer than 2 pairs of scales between the rostral and the first unpaired median head scale, ventral plates larger than dorsal scales, anterior nasals separated by rostral and frontonasal, and the lack of an ear opening.

• **DESCRIPTIONS.** In addition to original descriptions, detailed descriptions were provided by Burt and Burt (1931), Thomas (1965), Dixon (1973), Schwartz and Henderson (1991), and Vanzolini (1961a).

• **ILLUSTRATIONS.** Color photographs are in Powell et al. (1996), Murphy (1997), La Marca and Sori-ano (2004), Quinn et al. (2010), Hedges (2011), and Sánchez Muñoz (2011, including an image of an egg). Black-and-white photographs are in Ditmars (1928, 1933), Boos (1979), and Daudin and de Silva (2007, 2011). Line drawings are in Barbour (1914; dorsal, lateral, and ventral views of the head and views of the hand and foot), Beebe (1944; whole body, lateral view of head and forebody, and hindlimb), Brongersma (1946; lateral and ventral views of the head, forelimb, dorsal scales, and anal region of *B. anomala* [= *B. heteropa lineata*]), Vanzolini (1961a; 3 views of the head plus fore- and hindlimbs), Donoso-Barros and Garrido (1964; dorsal, lateral, and ventral views of the head, anal plates, dorsal and ventral scales, anterior and posterior limbs), Dixon (1973; prefrontal scales and hindlimb toes in the *B. heteropa* group in Venezuela), Ceí (2007; supraocular region); Thomas (1965), and Murphy (1997). Both of the latter illustrated dorsal head views of *B. h. allenii* and *B. h. trinitatis*. John et al. (2011) provided line drawings showing the use of the head in burrowing. Kohlsdorf and Wagner (2006) diagrammed digital loss in *B. h. trinitatis* and Galis et al. (2010) provided diagrammatic illustrations of gestalt and fore- and hindlimb morphology of *B. h. lineata* and reproduced the figure showing prefrontal scales and hindlimb toes in Dixon (1973). Presch (1975) illustrated fore- and hindlimb osteology, and the former was reproduced in Shapiro and Carl (2001) and Shapiro (2002). Ugeto and Rivas (2010) provided a color illustration of the entire animal, diagrammatic views of dorsal head squamation showing variation, and a color photograph of two preserved specimens.

• **DISTRIBUTION.** *Bachia heteropa* is found on the Grenada Bank (Bequia, Battowia, Mustique, Cannouan, Union, Grenada, and possibly Petit St. Vincent), Trinidad and satellites, Tobago, and through northern Venezuela (see also **Comment**). *Bachia heteropa* is mesophilic, at least semi-fossorial, and found in habitats with an abundance of ground litter (Barbour 1914; Dixon 1973; Ugeto and Rivas 2010). The distribution was previously illustrated by Dixon (1973). The West Indian range was illustrated in Schwartz and Henderson (1991), that for Isla de Margarita was mapped by Ugeto and Rivas (2010), and

distribution maps for *B. h. allenii* (Tobago) and *B. h. trinitatis* (Trinidad) are in Murphy (1992).

• **FOSSIL RECORD.** None.

• **PERTINENT LITERATURE.** The following literature citations pertaining to *B. heteropa* are arranged by topic: **behavior** (Donoso-Barros and Garrido 1964; John et al. 2011; Johnson 1946), **comparative morphology** (including limb loss) (Bergmann 2008; Ceí 2007; Dixon 1973; Galis et al. 2010; Kearney 2002; Kizirian and McDiarmid 1998; Kohlsdorf and Wagner 2006; Kohlsdorf et al. 2010; Lande 1978; MacLean 1974; Parker 1933; Presch 1975, 1980; Seligmann et al. 2003), **desiccation** (water loss rates) (Bentz et al. 2011), **diet** (Parker 1935), **distribution and zoogeography** (Daudin and de Silva 2007; Dixon 1973; Esqueda and La Marca 1999; Henderson and Powell 2006; Hoogmoed and Dixon 1977; Schwartz 1967), **geographic origin** (Hedges 1996; Murphy 1996), **key to subspecies** (Dixon 1973), **locomotion** (Beebe 1944; Donoso-Barros and Garrido 1964), **natural history** (Bentz et al. 2011; Boos 1979; Henderson and Powell 2009; Murphy 1997; Schwartz and Henderson 1991), **phylogenetics and systematics** (Dixon 1973; Galis et al. 2010; Kizirian and McDiarmid 1998; Kohlsdorf and Wagner 2006; Kohlsdorf et al. 2010), **predation** (Brongersma 1956; Murphy 1997; Ugeto and Rivas 2010), **tail autotomy** (Beebe 1944; Donoso-Barros and Garrido 1964), and **taxonomic history** (Brongersma 1946; Censky and Kaiser 1999; Peracca 1896; Thomas 1965).

The species was included under various synonyms in the following **checklists, keys, faunal accounts, or general works** (some include brief descriptions): Ayala (1986), Barbour (1914, 1930, 1935, 1937), Barbour and Loveridge (1929), Barros and Rincón (1999), Beebe (1952), Boos (1984a,b), Brygoo (1989), Burt and Burt (1933), Censky and Kaiser (1999), Cochran (1934), Cordero (1987), Corke (1992), Crother (1999), Daan and Hillenius (1966), Daudin and de Silva (2007), Donoso-Barros (1968), Esqueda et al. (2001), Evans (2008), Garman (1887), Germano et al. (2003), Gorzula and Señaris (1999), Groome (1970), Hailey and Cazabon-Mannette (2011), Hardy (1982), Henderson and Berg (2006, 2011), Henderson and Breuil (2012), Kramer (1979), La Marca and Soriano (2004), Lescure (1987), Lynn (1959), MacLean et al. (1977), Mahler and Kearney (2006), Malhotra and Thorpe (1999), Marcuzzi (1950), Markezich (2002), Mijares-Urrutia and Arends (1999, 2000), Mole and Urich (1894), Murphy (2004), Parker (1933, 1935), Pefaur (1992), Peters and Donoso-Barros (1970), Powell and Henderson (2005), Powell et al. (1996), Quinn et al. (2010), Rivas Fuenmayor et al. (2005), Romero Moreno and Romero Moreno (1989), Roux (1926), Ruthven (1925), Sánchez et al. (1995), Schwartz (1978), Schwartz and Henderson (1985, 1988), Schwartz and Thomas (1975), Shreve (1947), Ugeto and Rivas (2010), Underwood (1962), van Tuijl (1995), Vanzolini (1978, 1986), Werner (1900), and Wrobel (2005). Court (1858) mentioned “a curious little animal

resembling an amphisbaena, but having four limbs; it is probably a ceps.”

• **REMARKS.** Bauer and Günther (1994) noted that the original description mentioned 4 specimens, although measurements were given for only one. The ZMB catalogue lists 4 specimens under 3 consecutive numbers. All are apparently types. ZMB 1173, originally represented by 2 specimens listed as having been lost during World War II, is present, although only 1 specimen remains. The other members of the type-series originated from outside the currently construed range of *B. heteropa* (Dixon 1973), and their allocation to species should be further investigated. Authorship of *Chalcides heteropus* and other taxa first published in the “Nomenclator reptilium et amphibiorum Musei Zoologici Berolinensis” is here credited to Lichtenstein and von Martens. Although Lichtenstein alone is often given credit for names in this work, the preface clearly states that most of this work, including the descriptions, was due to von Martens (see Harris and Kluge 1984; Ulber 2003). David Weinland might also have played a role in the publication. Dixon (1973), Schwartz and Thomas (1975), Schwartz and Henderson (1988, 1991), and Powell et al. (1996) erroneously listed Natur-Museum Senckenberg (SMF) 39900 as the holotype. That specimen is the type of *Herpetochalcis heteropus* Boettger 1883 [*Bachia boettgeri* Boulenger 1885] (Mertens 1967). Smith and Taylor (1950) erroneously assumed that Boettger’s species was a synonym of *Chalcides heteropus*.

Lichtenstein (1856) indicated that the name was derived from a Wiegmann manuscript, explaining why Powell et al. (1996) listed the taxonomic authority as “Wiegmann in Lichtenstein.” However, as we are uncertain if Wiegmann also provided the description, authorship is most appropriately attributed to Lichtenstein and von Martens.

Dixon (1973), based on size and development of the limbs and toes, suggested that *B. h. lineata* and *B. h. marcelae* are predominately fossorial, *B. h. heteropa* is partially fossorial, and *B. h. alleni* and *B. h. trinitatis* are supraterranean leaf-litter inhabitants. Rivas Funemayor et al. (2005) called into question the validity of the various subspecies, citing clinal variation across the range of the species.

• **ETYMOLOGY.** The specific epithet is presumably from the Greek *heteros* (= different) and *pous* (= foot), almost certainly in reference to the variable size and number of hindlimb toes. The name *alleni* is a patronym honoring the collector of the holotype; *lineata* is from the Latin, presumably a reference to the striped pattern; the meaning of *marcelae* is unknown; and *trinitata* is derived from the type-locality (Trinidad) of this subspecies.

• **COMMENT.** *Chalcides Cuvieri* (Duméril and Bibron 1839) is included in synonymies of various subspecies of *B. heteropa* (e.g., Brongersma 1946, Dixon 1973). However, Brygoo (1989) clarified that the

specimens cited in Duméril and Bibron (1839) were from the banks of the Orinoco River and from Colombia, and are not attributable to *B. heteropa* (see also **Remarks**).

The Reptile Database (www.reptile-database.org) quoted F. Castro (pers. comm.) that the species has been found in Colombia. Ayala (1986) included *B. heteropa* in his list of Colombian lizards, based on a specimen from Norte de Santander in the Museo de Historia Natural, Instituto La Salle, Bogotá. Sánchez et al. (1995) listed Norte de Santander and Orinoquía as states of occurrence in Colombia.

1. *Bachia heteropa heteropa* (Lichtenstein and von Martens)

Chalcides heteropus Lichtenstein and von Martens 1856:17. See species synonymy.

Bachia heteropus: Garman 1892:97.

Bachia heteropa heteropa: Peters and Donoso-Barros 1970:80.

• **DEFINITION.** This subspecies is characterized (Dixon 1973) by tail squamation similar to body scales, prefrontals absent, supraoculars usually 2 (but sometimes 3 in intergrades between *B. h. heteropa* and *B. h. trinitatis*), 3 toes on hindlimbs, 7 rows of gular scales, 28–30 (mean 29.3) scales around midbody, 45–47 (mean 45.3) dorsal scale rows, and 27–30 (mean 28.3) ventral scale rows. One specimen examined by Dixon (1973) intermediate between *B. h. heteropa* and *B. h. trinitatis* had the second pair of chin shields in medial contact.

The 3 dorsal dark lines are usually evident, occasionally irregular, with the light interspatial stripes lightly freckled or mottled (Dixon 1973).

2. *Bachia heteropa alleni* (Barbour)

Scolecocaurus cuvieri: Boulenger 1885:416 (part).

Scolecocaurus alleni Barbour 1914:315. Type-locality, “St. George’s, Grenada.” Holotype, Museum of Comparative Zoology (MCZ) 7793, an adult, sex unknown, collected by G.M. Allen, date of collection unknown (not examined by authors). *Lapsus*.

Scolecocaurus alleni alleni: Barbour 1933:76.

Scolecocaurus alleni parviceps Barbour 1933:77. Type-locality, “Cannouan, The Grenadine Islands, B.W.I.” Holotype, Museum of Comparative Zoology (MCZ) 32345, an adult, sex unknown, collected by D. Fairchild and H.F. Loomis on 6 February 1932 (not examined by authors).

[*Bachia*] *alleni*: Vanzolini 1961b:204.

Scolecocaurus trinitatis: Underwood 1962:94 (part).

Scolecocaurus trinitatis parviceps: Underwood 1962:94.

Bachia alleni alleni: Thomas 1965:145.

Scolecocaurus sp.: Groome 1970:36.

Bachia heteropa alleni: Dixon 1973:32.

Bachia heteropus alleni: Schwartz and Thomas 1975:109.

B[achia]. (*Scolecocaurus*) *alleni*: Obst, Richter, and Jacob 1984:91.

• **DEFINITION.** This subspecies is characterized (Dixon 1973) by tail squamation mostly hexagonal, prefrontals present and in contact medially, 3 supraoculars, 4 toes on hindlimbs, 8 rows, of gular scales, 27–31 (mean 29.0) scale rows around midbody, 39–45 (mean 40.9) dorsal scale rows, and 24–29 (mean 26.4) ventral scale rows.

Coloration varies slightly from the basic pattern by tending towards an obscure median dark line and paravertebral dark lines broken into a series of dashes or spots with the light interspaces heavily mottled or freckled with dark brown (Dixon 1973).

3. *Bachia heteropa lineata* (Boulenger)

Bachia lineata Boulenger 1903:432. Type-locality, “Duaca, Venezuela.” Holotype, British Museum (Natural History) (BMNH) 1946.8.2.41 (formerly 1902.7.29.90), an adult (sex unknown), collected by Mr. Wayman (purchased from Mr. Rosenberg), date of collection unknown, but registered into the collections on 29 July 1902 (examined by C. McCarthy at the request of the authors).

Bachia anomala Roux 1929:31. Type-locality, “El Mene, Distr. Acosta, Prov. Falcon, Vénézuéla.” Holotype, Naturhistorisches Museum, Basel (NMBA) 9912, an adult male (SVL 52 mm), collected by H.G. Kugler and presented to the museum in 1929 (probably collected in the same year; R. Winkler, in litt., 29.XI.2010) (not examined by authors).

Bachia heteropa lineata: Mijares-Urrutia and Arends 1999:115.

• **DEFINITION.** This subspecies is characterized by tail squamation similar to body scales, prefrontals absent, 2 supraoculars, 2 toes on hindlimbs, 7 rows of gular scales, 24–26 (mean 25.3) scales around midbody, 45–49 (mean 47.2) dorsal scale rows, and 24–26 (mean 25.3) ventral scale rows (Dixon 1973).

The 3 dorsal dark lines are usually evident, occasionally irregular, with the light interspatial stripes lightly freckled or mottled (Dixon 1973).

• **REMARKS.** Shreve (1947) suggested that *B. lineata* and *B. anomala* might not be conspecific.

4. *Bachia heteropa marcelae* Donoso-Barros and Garrido

Bachia marcelae Donoso-Barros and Garrido 1964:1. Type-locality, “Bosque tropical La Luz, Edo. Barinas, Venezuela.” Holotype, D-B 6312061, collected on 12 December 1963 by R. Donoso-Barros and A. Garrido (not examined by authors).

Bachia lineata marcelae: Donoso-Barros 1968:117.

Bachia heteropa marcelae: Dixon 1973:33.

• **DEFINITION.** This subspecies is characterized by tail squamation similar to body scales, prefrontals absent, 2 supraoculars, 2 toes on hindlimbs, 5 rows of gular scales, 25 (in 3 specimens) scale rows around midbody, and 39–42 (mean 40.7) dorsal scale rows (Dixon 1973).

The 3 dorsal dark lines are usually evident, occasionally irregular, with the light interspatial stripes lightly freckled or mottled (Dixon 1973).

5. *Bachia heteropa trinitatis* (Barbour)

Scolecocaurus cuvieri: Boettger 1893:76.

Scolecocaurus trinitatis Barbour 1914:316. Type-locality, “Caparo, Trinidad.” Holotype, Museum of Comparative Zoology (MCZ) 8947, an adult, sex unknown, collected by A.B. Carr, date of collection unknown (not examined by authors).

[*Bachia*] *trinitatis*: Vanzolini 1961b:204.

Bachia alleni trinitatis: Thomas 1965:152.

Bachia heteropa trinitatis: Dixon 1973:32.

Bachia heteropa trinitatus: MacLean, Kellner, and Dennis 1977:45. *Lapsus*.

Bachia trinitatis: Kenny 1979:9.

• **DEFINITION.** This subspecies is characterized (Dixon 1973) by tail squamation mostly hexagonal, prefrontals present (not in contact medially), 3 supraoculars, 4 toes on hindlimb, 7 rows of gular scales in 57.5% of specimens (8 rows of gular scales in the rest), 26–31 (mean 28.5) scales around midbody, and 38–45 (mean 41.3) dorsal scale rows.

The basic color pattern is retained in this subspecies with 3 dark lines, but these can be broken or irregular. The 2 paravertebral light interspatial stripes usually are prickled with dark brown, whereas the 2 dorsolateral light interspatial stripes are uniform tan to yellowish brown (Dixon 1973).

• **ACKNOWLEDGEMENTS.** Colin McCarthy (BMNH), José Rosado (MCZ), and Raffael Winkler (NMBA) examined type-specimens in their care. Adrian Hailey, John C. Murphy, and Gilson Rivas helped locate obscure references. John S. Parmelee, Jr. prepared the map.

LITERATURE CITED

- Ávila Pires, T.C.S. de. 2005. Reptiles, p. 25–40. *In* T. Hollowell and R.P. Reynolds (eds.), Checklist of the terrestrial vertebrates of the Guiana Shield. Bull. Biol. Soc. Washington (13).
- Ayala, S. 1986. Saurios de Colombia: Lista actualizada y distribución de ejemplares colombianos en los museos. *Caldasia* 15:555–575.
- Barbour, T. 1914. A contribution to the zoögeography of the West Indies, with special reference to amphibians and reptiles. *Mem. Mus. Comp. Zool.* 44: 205–359.
- . 1930. A list of Antillean reptiles and amphibians. *Zoologica* (N.Y.) 11:61–116.
- . 1933. Notes on *Scolecocaurus*. *Copeia* 1933:74–77.
- . 1935. A second list of Antillean reptiles and amphibians. *Zoologica* (N.Y.) 19:77–141.
- . 1937. Third list of Antillean reptiles and amphibians. *Bull. Mus. Comp. Zool.* 82:77–166.
- Barros, T. and A.D. Rincón R. 1999. Segundo registro y redescipción de *Bachia bicolor* (Sauria: Gymnophthalmidae) en Venezuela. *Bol. Centro Invest.*

- Biol. 33:111–124.
- Bauer, A.M. and R. Günther 1994. An annotated type catalogue of the teiid and microteiid lizards in the Zoological Museum, Berlin (Reptilia: Squamata: Teiidae and Gymnophthalmidae). Mitt. Zool. Mus. Berlin 70:267–280.
- Beebe, W. 1945. Field notes on the lizards of Kartabo, British Guiana, and Caripito, Venezuela. Part 3. Teiidae, Amphisbaenidae and Scincidae. Zoológica (N.Y.) 30:7–32, pls. I–V.
- . 1952. Introduction to the ecology of the Arima Valley, Trinidad, B.W.I. Zoologica (N.Y.) 37:157–183 + 5 pls.
- Bentz, E.J., M.J. Rivera Rodríguez, R.R. John, R.W. Henderson, and R. Powell. 2011. Population densities, activity, microhabitats, and thermal biology of a unique crevice and litter-dwelling assemblage of reptiles on Union Island, St. Vincent and the Grenadines. Herpetol. Conserv. Biol. 6:40–50.
- Bergmann, P.J. 2008. A Phylogenetic Study and Functional Approach to the Study of the Evolution of Body Shape in Lizards (Squamata). Ph.D. Diss., Univ. Massachusetts, Amherst.
- Boettger, O. 1883. Herpetologische Mittheilungen. I. Kurze Notizen über Reptilien und Amphibien in der Heidelberger Universitäts-Sammlung. Ber. Offenbach. Verh. Naturk. 22/23:147–152.
- . 1893. Katalog der Reptilien-Sammlung im Museum der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main. I. Teil (Rhynchocephalen, Schildkröten, Krokodile, Eidechsen, Chamäleons). Gebrüder Knauer, Frankfurt a. M.
- Boos, H.E.A. 1979. Teiids part 2. Trinidad Nat. 2(7): 23.
- . 1984a. A consideration of the terrestrial reptile fauna on some offshore islands north west of Trinidad. Living World. J. Trinidad Tobago Fld. Nat. Club. 1983–1984:19–26.
- . 1984b. The terrestrial reptiles of Monos Island. Living World. J. Trinidad Tobago Fld. Nat. Club. 1983–1984:14–18.
- Boulenger, G.A. 1885. Catalogue of the lizards in the British Museum (Natural History). Vol. II. Trustees (Brit. Mus.), London.
- . 1903. Descriptions of new lizards in the collection of the British Museum. Ann. Mag. Nat. Hist. ser. 7, 12:429–435.
- Brongersma, L.D. 1946. Some notes on species of the genera *Bachia* and *Scolecocaurus*. Zool. Meded. Leiden 26:237–246.
- . 1956. On some reptiles and amphibians from Trinidad and Tobago, B.W.I. II. Proc. Konink. Nederland Akad. Wetensch., Ser. C, 59(2):177–188.
- Brygoo, E.R. 1989. Les types de teiids (Reptiles, Sauriens) du Muséum national d'Histoire naturelle, catalogue critique. Bull. Mus. Nat. d'Hist. Nat. (4) 11A (no 1, suppl.):1–44.
- Burt, C.E. and M.D. Burt. 1931. South American lizards in the collection of the American Museum of Natural History. Bull. Amer. Mus. Nat. Hist. 61: 227–395.
- and –. 1933. A preliminary check list of the lizards of South America. Trans. Acad. Sci. St. Louis 28: 1–104.
- Cei, J.M. 2007. Comparative studies of supraocular lepidosis in Squamata (Reptilia) and its relationships with an evolutionary taxonomy. Multequina (16):1–52.
- Censky, E.J. and H. Kaiser. 1998. Lesser Antillean Herpetofauna. In B.I. Crother (ed.), Caribbean Amphibians and Reptiles. Academic Press, San Diego, California.
- Cochran, D.M. 1934. Herpetological collections from the West Indies made by Dr. Paul Bartsch under the Walter Rathbone Bacon Scholarship, 1928–1930. Smithsonian Misc. Coll. 92:1–48.
- Cordero, G.A. 1987. Composición y diversidad de la fauna de vertebrados terrestres de Barlovento, Estado Miranda, Venezuela. Acta Cient. Venez. 38: 234–258.
- Corke, D. 1992. The status and conservation needs of the terrestrial herpetofauna of the Windward Islands (West Indies). Biol. Conserv. 62:47–58.
- Court, J. 1858. Catalogue of reptiles, p. 440–446. In L.A.A. Verteuil, Trinidad: Its Geography, Natural Resources, Administration, Present Condition, and Prospects. Ward and Lock, London.
- Crother, B.I. 1999. Evolutionary relationships, p. 269–334. In B.I. Crother (ed.), Caribbean Amphibians and Reptiles. Academic Press, San Diego, California.
- Daan, S. and D. Hillenius. 1966. Catalogue of the type specimens of amphibians and reptiles in the Zoological Museum, Amsterdam. Beaufortia 13: 117–144.
- Daudin, J. and M. de Silva. 2007. An annotated checklist of the amphibians and terrestrial reptiles of the Grenadines with notes on their local natural history and conservation. Appl. Herpetol. 4:163–175.
- and –. 2011. An annotated checklist of the amphibians and terrestrial reptiles of the Grenadines with notes on their local natural history and conservation, p. 259–272. In A. Hailey, B.S. Wilson, and J.A. Horrocks (eds.), Conservation of Caribbean Island Herpetofaunas. Vol. 2: Regional Accounts of the West Indies. Brill, Leiden, The Netherlands.
- Ditmars, R.L. 1928. Reptiles of the World. Macmillan, New York.
- . 1933. Reptiles of the World. Rev. ed. Macmillan, New York.
- Dixon, J.R. 1973. A systematic review of the teiid lizards, genus *Bachia*, with remarks on *Heterodactylus* and *Anotosaurus*. Misc. Publ. Univ. Kansas Mus. Nat. Hist. (57):1–47.
- Donoso-Barros, R. 1968. The lizards of Venezuela (check list and key). Carib. J. Sci. 8:105–122.
- and A. Garrido R. 1964. Nuevo Teiidae de Venezuela *Bachia marcelae*, nov. sp. Publ. Ocas. Mus. Cienc. Nat., Caracas (Zool.) 8:1–7.
- Duméril, A.M.C. and G. Bibron 1839. Erpétologie Générale ou Histoire Naturelle des Reptiles. Vol. 5. Libr. Encyclopedique de Roret, Paris.
- Esqueda, L.F. and E. La Marca. 1999. New reptilian species records from the Cordillera de Mérida, Andes of Venezuela. Herpetol. Rev. 30:238–240.

- , E. la Marca, M. Natera, and P. Battiston. 2001. Noteworthy reptilian state records and a lizard species new to the herpetofauna of Venezuela. *Herpetol. Rev.* 32:198–200.
- Evans, S.E. 2008. The skull of lizards and tuatara, p. 1–347. *In* C. Gans, A.S. Gaunt, and K. Adler (eds.), *Biology of the Reptilia*, Vol. 20, Morphology H. Society for the Study of Amphibians and Reptiles, Ithaca, New York.
- Fitzinger, L.J. 1826. Neue Classification der Reptilien nach ihren natürlichen Verwandtschaften, nebst einer Verwandtschafts-Tafel und einem Verzeichnisse der Reptilien-Sammlung des K.K. Zoologischen Museum's zu Wien. J.G. Heubner, Wien. [8] + 66 pp., 1 foldout table.
- Galis, F., J.W. Arntzen, and R. Lande. 2010. Dollo's law and the irreversibility of digit loss in *Bachia*. *Evolution* 64:2466–2476.
- Garman, S. 1887. On the West Indian Teiidae in the Museum of Comparative Zoology. *Bull. Essex Inst.* 19:1–12.
- . 1892. On *Cophias* and *Bachia*. *Bull. Essex Inst.* 24: 96–97.
- Germano, J.M., J.M. Sander, R.W. Henderson, and R. Powell. 2003. Herpetofaunal communities in Grenada: A comparison of altered sites, with an annotated checklist of Grenadian amphibians and reptiles. *Carib. J. Sci.* 39:68–76.
- Gorzula, S. and J.C. Señaris. 1999. Contribution to the herpetofauna of the Venezuelan Guayana. I. A Data Base. *Sci. Guaianae* (8):xviii + 269 p., 129 photographs, 4 maps.
- Groome, J.R. 1970. A Natural History of the Island of Grenada, West Indies. *Carib. Printers, Ltd., Arima, Trinidad*.
- Hailey, A. and M. Cazabon-Mannette. 2011. Conservation of herpetofauna in the Republic of Trinidad and Tobago, p. 183–217. *In* A. Hailey, B.S. Wilson, and J.A. Horrocks (eds.), *Conservation of Caribbean Island Herpetofaunas*. Vol. 1: Conservation Biology and the Wider Caribbean. Brill, Leiden, The Netherlands.
- Hardy, J.D. 1982. Biogeography of Tobago, West Indies, with special reference to amphibians and reptiles: A review. *Bull. Maryland Herpetol. Soc.* 18:37–142.
- Harris, D.M. and A.G. Kluge. 1984. The *Sphaerodactylus* (Sauria: Gekkonidae) of Middle America. *Occas. Pap. Mus. Zool. Univ. Michigan* (706):1–59.
- Hedges, S.B. 1996. The origin of West Indian amphibians and reptiles, p. 95–128. *In* R. Powell and R.W. Henderson (eds.), *Contributions to West Indian Herpetology: A Tribute to Albert Schwartz*. *Contrib. Herpetol. Soc. Study Amphib. Rept.* (12).
- Hedges, S.B. 2011. *CaribHerp*. <<http://caribherp.org/>>.
- Henderson, R.W. and C.S. Berg. 2006. The herpetofauna of Grenada and the Grenada Grenadines: Conservation concerns. *Appl. Herpetol.* 3:197–213.
- and –. 2011. The herpetofauna of Grenada and the Grenada Grenadines: Conservation concerns, p. 239–258. *In* A. Hailey, B.S. Wilson, and J.A. Horrocks (eds.), *Conservation of Caribbean Island Herpetofaunas*. Vol. 2: Regional Accounts of the West Indies. Brill, Leiden, The Netherlands.
- and M. Breuil. 2012. Lesser Antilles, p. 150–161. *In* R. Powell and R.W. Henderson (eds.), *Island lists of West Indian amphibians and reptiles*. *Bull. Florida Mus. Nat. Hist.* 51:87–168.
- and R. Powell. 1999. West Indian herpetoecology, p. 223–268. *In* B.I. Crother (ed.), *Caribbean Amphibians and Reptiles*. Academic Press, San Diego, California.
- and –. 2006. Geographic distribution: *Bachia heteropa alleni*. *Herpetol. Rev.* 37:360.
- and –. 2009. Natural History of West Indian Reptiles and Amphibians. Univ. Press Florida, Gainesville.
- Hoogmoed, M.S. and J.R. Dixon. 1977. A new species of *Bachia* (Teiidae, Sauria) from Estado Bolívar, Venezuela, with notes on the zoogeography of the genus. *Zool. Meded.* 51:25–31.
- John, R.R., E.J. Bentz, M.J. Rivera Rodríguez, M.E. Gifford, and R. Powell. 2011. *Bachia heteropa alleni* (Earless Worm Lizard). Escape and digging behaviors. *Herpetol. Rev.* 42:92.
- Johnson, M.L. 1946. Herpetological notes from Trinidad. *Copeia* 1946:108.
- Kearney, M. 2002. Appendicular skeleton in amphisbaenians (Reptilia: Squamata). *Copeia* 2002: 719–738.
- Kenny, J.S. 1979. Floorplan, environment and fauna of Tamana caves. *Living World, J. Trinidad Tobago Fld. Nat. Club* 1978–1979:5–9.
- Kizirian, D.A. and R.W. McDiarmid. 1998. A new species of *Bachia* (Squamata: Gymnophthalmidae) with plesiomorphic limb morphology. *Herpetologica* 54:245–253.
- Kohlsdorf, T. and G.P. Wagner. 2006. Evidence for the reversibility of digit loss: A phylogenetic study of limb evolution in *Bachia* (Gymnophthalmidae: Squamata). *Evolution* 60:1896–1912.
- , V.J. Lynch, M.T. Rodrigues, M.C. Brandley, and G.P. Wagner. 2010. Data and data interpretation in the study of limb evolution: A reply to Galis et al. on the revolution of digits in the lizard genus *Bachia*. *Evolution* 64:2477–2485.
- La Marca, E., J. Manzanilla, and A. Mijares-Urrutia. 2004. Revisión taxonómica del *Colostethus* de norte de Venezuela confundido durante largo tiempo con *C. brunneus*. *Herpetotropicos* 1(4): 40–50.
- and P.J. Soriano. 2004. Reptiles de Los Andes de Venezuela. Catálogo Zoológico de Venezuela, Volumen 2. BIOGEOS, Mérida, Venezuela.
- Lande, R. 1978. Evolutionary mechanisms of limb loss in tetrapods. *Evolution* 32:73–92.
- Lescure, J. 1987. Le peuplement en reptiles et amphibians des Petites Antilles. *Bull. Soc. Zool. France* 112:327–342.
- Lichtenstein, H. 1856. Nomenclator reptilium et amphibiorum Musei Zoologici Berolinensis. Namenverzeichnis der in der zoologischen Sammlung der Königlichen Universität zu Berlin aufgestellten Arten von Reptilien und Amphibien nach ihren

- Ordnungen, Familien und Gattungen. Königl. Akad. Wissensch., Berlin.
- Lynn, W.G. 1959. Some Reptiles and Amphibians from Trinidad. *Herpetologica* 15:113–117.
- MacLean, W.P. 1974. Feeding and locomotor mechanisms of teiid lizards: Functional morphology and evolution. *Pap. Avul. Zool. S. Paulo* 27:179–213.
- , R. Kellner, and H. Dennis. 1977. Island lists of West Indian amphibians and reptiles. *Smithson. Herpetol. Info. Serv.* (40):1–47.
- Mahler, L. and M. Kearney. 2006. The palatal dentition in squamate reptiles: Morphology, development, attachment, and replacement. *Fieldiana Zool.*, n. s. (108):1–61.
- Malhotra, A. and R.S. Thorpe. 1999. Reptiles and Amphibians of the Eastern Caribbean. MacMillan Educ. Ltd., London.
- Marcuzzi, G. 1950. Breves apuntes sobre algunos lagartos de Venezuela Septentrional. *Mem. Soc. Cien. Nat. La Salle* 10:73–110.
- Markezich, AL. 2002. New distribution records of reptiles from western Venezuela. *Herpetol. Rev.* 33: 69–74.
- Mertens, R. 1967. Die herpetologische Sektion des Natur-Museums und Forschungs-Institutes Senckenberg in Frankfurt a. M. nebst einem Verzeichnis ihrer Typen. 1. *Senckenberg. Biol.* 48:1–106 + 4 pls.
- Mijares-Urrutia, A. and A. Arends R. 1999. Additional new regional and local records of amphibians and reptiles from the state of Falcón, Venezuela. *Herpetol. Rev.* 30:115.
- and –. 2000. Herpetofauna of Estado Falcón, north-western Venezuela: a checklist with geographical and ecological data. *Smithson. Herpetol. Info. Serv.* (123):1–30.
- Mole, R.R. and F.W. Urich. 1894. A preliminary list of the reptiles and batrachians of the Island of Trinidad. *Living World, J. Trinidad Fld. Nat. Club* 2: 77–90.
- Murphy, J.C. 1996. Crossing Bond's Line: The herpetofaunal exchange between the eastern Caribbean and mainland South America, p. 207–216. *In* R. Powell and R.W. Henderson (eds.), *Contributions to West Indian Herpetology: A Tribute to Albert Schwartz*. Soc. Study Amphib. Rept., Contrib. Herpetol. (12).
- . 1997. Amphibians and Reptiles of Trinidad and Tobago. Krieger Publ. Co., Malabar, Florida.
- Murphy, W.L. 2004. A Birdwatchers' Guide to Trinidad and Tobago. Prion Ltd., Cley next the Sea, United Kingdom.
- Obst, F.J., K. Richter, and U. Jacob. 1984. *Lexikon der Terraristik und Herpetologie*. Ed. Leipzig, Leipzig, Germany (English translation: 1988. *The Completely Illustrated Atlas of Reptiles and Amphibians for the Terrarium*. T.F.H. Publ., Inc., Neptune City, New Jersey).
- Parker, H.W. 1933. Some amphibians and reptiles from the Lesser Antilles. *Ann. Mag. Nat. Hist. Ser.* 10:151–158.
- . 1935. The lizards of Trinidad. *Trop. Agric.* 12:65–70.
- Pefaur, J.E. 1992. Checklist and bibliography (1960–85) of the Venezuelan herpetofauna. *Smithson. Herpetol. Info. Serv.* (89):1–54.
- Peracca, M.G. 1896. Rettili ed anfibi raccolti nel Darien ed a Panama dal dott. E. Festa. *Boll. Mus. Zool. Anat. Comp. Torino* 253:1–12.
- Peters, J.A. and R. Donoso-Barros. 1970. Catalogue of the Neotropical Squamata: Part II Lizards and Amphisbaenians. *Smithson. Inst. Press*, Washington, D.C.
- Powell, R. and R.W. Henderson. 2005. Conservation status of Lesser Antillean reptiles. *Iguana* 12:62–77.
- , –, K. Adler, and H.A. Dundee. 1996. A checklist of West Indian amphibians and reptiles, p. 51–93 + 8 pl. *In* R. Powell and R.W. Henderson (eds.), *Contributions to West Indian Herpetology: A Tribute to Albert Schwartz*. Contrib. Herpetol. Soc. Study Amphib. Rept. (12).
- Presch, W. 1975. The evolution of limb reduction in the teiid lizard genus *Bachia*. *Bull. So. California Acad. Sci.* 74:113–121.
- . 1980. Evolutionary history of the South American microteiid lizards (Teiidae: Gymnophthalmine). *Copeia* 1980:36–56.
- Quinn, D.P., A.L. McTaggart, T.A. Bellah, E.J. Bentz, L.G. Chambers, H.D. Hedman, R.R. John, D.N. Muñiz Pagan, and M.J. Rivera Rodriguez. 2010. The reptiles of Union Island, St. Vincent and the Grenadines. *Rept. & Amphib.* 17:222–233.
- Rivas Fuenmayor, G., G. Ugueto, R. Rivero, and A. Miralles. 2005. The herpetofauna of Isla de Margarita, Venezuela: New records and comments. *Carib. J. Sci.* 41:346–351.
- Romero Moreno, M.E. and C. Romero Moreno. 1989. Desde el Orinoco hacia el Siglo XXI: El Hombre, La Fauna y Su Medio. Fondo FEN Colombia, Bogotá, Colombia.
- Roux, J. 1926. Notes d'erpétologie sud-américaine. *Rev. Suisse Zool.* 33:291–299.
- . 1929. Sur deux reptiles nouveaux du Venezuela. *Verh. Nat. Ges. Basel* 40:29–34.
- Ruthven, A.G. 1925. Lizards of the genus *Bachia*. *Proc. Boston Soc. Nat. Hist.* 38:101–109.
- Sánchez-C., H., O.M. Castaño-M., and G. Cárdenas-A. 1995. Diversidad de los reptiles en Colombia, p. 277–325. *In* J.O. Rangel (ed.), *Colombia: Diversidad Biótica*, Vol. 1. Editorial Guadalupe, Santa Fé de Bogotá, Colombia.
- Sánchez Muñoz, A.J. 2011. Father Sanchez's Web Site of West Indian Natural History. <www.kingsnake.com/westindian/>.
- Schwartz, A. 1967. Frogs of the genus *Eleutherodactylus* in the Lesser Antilles. *Stud. Fauna Curaçao Carib. Isl.* (91):1–62.
- . 1978. Some aspects of the herpetogeography of the West Indies, p. 31–51. *In* F.B. Gill (ed.), *Zoogeography in the Caribbean*. The 1975 Leidy Medal Symposium. Acad. Nat. Sci. Philadelphia Spec. Publ. (13).
- and R.W. Henderson. 1985. A Guide to the Identification of the Amphibians and Reptiles of the West Indies Exclusive of Hispaniola. Milwaukee Pub.

- Mus., Milwaukee, Wisconsin.
- and –. 1988. West Indian amphibians and reptiles: a checklist. *Contrib. Biol. Geol. Milwaukee Pub. Mus.* (74):1–264.
 - and –. 1991. *Amphibians and Reptiles of the West Indies: Descriptions, Distributions, and Natural History*. Univ. Florida Press, Gainesville.
 - and R. Thomas. 1975. A check-list of West Indian amphibians and reptiles. *Carnegie Mus. Nat. Hist. Spec. Publ.* (1):1–216.
- Seligmann, H., A. Beiles, and Y.L. Werner. 2003. More injuries in left-footed individual lizards and *Sphenodon*. *J. Zool., Lond.* 260:129–144.
- Shapiro, M.D. 2002. Developmental morphology of limb reduction in *Hemiergis* (Squamata: Scincidae): Chondrogenesis, osteogenesis, and heterochrony. *J. Morphol.* 254:211–231.
- and T.F. Carl. 2001. Novel features of tetrapod limb development in two non-traditional model species: A skink and a direct-developing frog, p. 337–361. *In* M. Zelditch (ed.), *Beyond Heterochrony: The Evolution of Development*. Wiley-Liss, New York.
- Shreve, B. 1947. On Venezuelan reptiles and amphibians collected by Dr. G. Kugler. *Bull. Mus. Comp. Zool.* 99:517–537.
- Smith, H.M. and E.H. Taylor. 1950. An annotated checklist and key to the reptiles of Mexico, exclusive of the snakes. *Bull. U.S. Natl. Mus.* (199):v + 253 p.
- Thomas, R. 1965. The smaller teiid lizards (*Gymnophthalmus* and *Bachia*) of the southeastern Caribbean. *Proc. Biol. Soc. Washington* 78:141–154.
- Ulber, T. 2003. Annotated index, p. iv–v. *In* T. Ulber (ed.), *Rare Herpetological Literature I*. Herprint Intl., Bredell.
- Ugeto, G.N. and G.A. Rivas. 2010. *Amphibians and Reptiles of Margarita, Coche and Cubagua*. Ed. Chimaira, Frankfurt am Main.
- Underwood, G. 1962. Reptiles of the eastern Caribbean. *Carib. Affairs* (n.s.) 1:1–191.
- van Tuijl, L. 1995. Revised catalogue of the type specimens of recent amphibians and reptiles in the “Zoologisch Museum” University of Amsterdam, the Netherlands. *Bull. Zoöl. Mus.* 14:125–144.
- Vanzolini, P.E. 1961a. Redescricao de *Scolecosaurius trinitatis* (Sauria, Teiidae). *Pap. Avul. Zool. S. Paulo* 14:183–185.
- . 1961b. *Bachia*: espécies brasileiras e conceito genérico (Sauria, Teiidae). *Pap. Avul. Zool. S. Paulo* 14:193–209.
 - . 1978. An Annotated Bibliography of the Land and Fresh-water Reptiles of South America (1958–1975). Vol. II (1901–1975). *Mus. Zool., Univ. São Paulo, São Paulo, Brazil*.
 - . 1986. Addenda and corrigenda to part II lizards and Amphisbaenia, p. 1–23. *In* J.A. Peters and R. Donoso-Barros, *Catalogue of the Neotropical Squamata: Part II Lizards and Amphisbaenians*, rev. ed. *Smithson. Inst. Press, Washington, D.C.*
- Wagler, J. 1830. *Natürliches System der Amphibien mit vorangehender Classification der Säugthiere und Vögel. Ein Beitrag zur vergleichenden Zoologie*. J. G. Cotta'schen Buchhandlung, München, Stuttgart, Tübingen.
- Werner, F. 1900. Ueber Reptilien und Batrachier aus Colombien und Trinidad, II. *Verh. Zool.-Bot. Ges. Wien* 50:262–272.
- Wrobel, M. 2005. *Elsevier's Dictionary of Reptiles*. Elsevier, Amsterdam.

Rebecca R. John, Departments of Environmental Studies and Biology, University of California, Santa Cruz, CA 95064, USA (lizardgeeky@yahoo.com), **Mel José Rivera Rodríguez**, Department of Biology, University of Puerto Rico, Mayagüez, PR 00681 (mel-rivera103@yahoo.com), **Ehren J. Bentz**, Feather River College, Quincy, CA 95971, USA (Current address: Department of Biology, Oregon State University, Corvallis, OR 97331, USA) (ehren_bentz@yahoo.com), **Aaron M. Bauer**, Department of Biology, Villanova University, Villanova, PA 19085, USA (aaron.bauer@villanova.edu), and **Robert Powell**, Department of Natural Sciences, Avila University, Kansas City, MO 64145, USA (robert.powell@avila.edu).

Primary editor for this account, Andrew H. Price.

Published 30 April 2012 and Copyright © 2012 by the Society for the Study of Amphibians and Reptiles.
